

TABLE OF CONTENTS

EXECUTIVE SUMMARY		
1	INTRODUCTION.....	1
2	ALGAL CULTURE - A RED HERRING?.....	7
	2.1 Introduction.....	7
	2.2 Water use and yields.....	8
	2.2.1 Extensively in ponds.....	8
	2.2.2 Intensively in ponds	8
	2.2.3 Confined in existing water bodies	10
	2.3 Water use in aquaculture vs irrigated crop production	11
	2.4 Protein requirements of South Africa	17
	2.5 Quantity of agricultural waste available	18
	2.6 Algal production values	20
	2.7 Conversion of waste to biomass	23
	2.8 Conversion of algal protein to animal protein	24
	2.9 Candidate species for algalculture	27
	2.10 Conclusion	28
3	METHODS AND MATERIALS	29
	3.1 The release of nutrients from animal waste	29
	3.1.1 Source of waste	29
	3.1.2 Waste loss during fermentation	29
	3.1.3 Nutrient release from anaerobic ferment	29
	3.1.4 Nutrient requirements of <i>Spirulina</i>	30
	3.2 Method of aeration	30
	3.2.1 Intermittent aeration	30
	3.2.2 Continuous aeration	30
	(i) Unbuffered pH	30
	(ii) Buffered pH	31
	3.3 The tolerance of <i>Spirulina</i> to NaCl, Ca ²⁺ and SO ₄ ²⁻	31
	3.4 The influence of pH on rotifers	32
	3.5 Definition of terms	32
	3.6 Use of the fairy shrimp <i>Streptocephalus macrourus</i> for the recovery of nutrients in waste water	32
	3.6.1 Batch cultures	32
	3.6.2 Continuous cultures	33
	3.6.3 Estimation of dry mass	33
	3.6.4 Calculation of growth efficiency	33
	3.6.5 Calculation of daily growth rate	34
	3.6.6 Estimation of production	34

4	RESULTS AND DISCUSSION	35
4.1	The release of nutrients from animal waste	35
4.1.1	Introduction	35
4.1.2	Waste loss during fermentation	35
4.1.3	Nutrient release from anaerobic ferment	35
4.1.4	Nutrient requirements of <i>Spirulina</i>	39
4.1.5	Discussion	39
4.2	Method of aeration	39
4.2.1	Intermittent aeration	39
4.2.2	Continuous aeration	40
4.3	The tolerance of <i>Spirulina</i> to NaCl, Ca ²⁺ and SO ₄ ²⁻	40
4.3.1	NaCl	41
4.3.2	Ca ²⁺	41
4.3.3	SO ₄ ²⁻	44
4.4	The influence of pH on rotifers	46
4.4.1	Unbuffered pH	46
4.4.2	Controlled pH	49
4.5	Use of the fairy shrimp <i>Streptocephalus macrourus</i> for the recovery of the nutrients in waste water	60
4.5.1	Batch culture	62
4.5.2	Growth efficiency	62
4.5.3	Production	65
4.5.4	Continuous cultures	67
4.5.5	Growth efficiencies of <i>S. macrourus</i> in mass culture under different management conditions	71
4.5.6	Assessment of <i>S. macrourus</i> as an aquaculture candidate	74
5	THE ECONOMICS OF <i>Spirulina</i> PRODUCTION ON AGRICULTURAL WASTES FOR USE AS A STOCK FEED	76
5.1	Harvesting of algae	76
5.2	Cost estimate for the production of biomass	77
5.3	Economics of the production unit	80
6	CONCLUSIONS	84
7	ACKNOWLEDGEMENTS	86
8	REFERENCES	87