TABLE OF CONTENTS

Executive summary	**i**
Acknowledgements	**ix**
List of Tables	**xvi**
List of Figures	**xix**
List of Acronyms and Abbreviations	**xxiii**
CHAPTER 1: INTRODUCTION	**1**
1.1 SCOPE	1
1.2 OBJECTIVES	2
1.3 APPROACH	2
1.3.1 Survey to determine the extent to which sacrificial land disposal is used in SA	2
1.3.2 Detailed study of selected sites	3
1.3.3 Estimation of the potential future impact of dedicated land disposal of sewage sludge on the water environment	4
CHAPTER 2: LITERATURE REVIEW	**5**
2.1 The problem	5
2.2 Dedicated land disposal (DLD)	6
2.3 The advantages and disadvantages of DLD	8
2.4 The potential pollution problem	8
2.5 International approach to sludge disposal	9
2.5.1 Australia	10
2.5.2 Canada	11
2.5.3 Europe	12
2.5.4 United Kingdom	12
2.5.5 USA	13
2.6 Guidelines for potentially toxic metals and elements (PTME) in soil and water	**14**
2.7 Modelling of metal migration in sludge ammended soils	18

2.7.1 Metal Contaminant Mobility and bioavailability in soils 18

2.7.2 Metal ion complexing capacity in sewage-associated water 18

2.7.3 Movements of metals within soil profiles 19

2.7.3.1 Effect of pH 20

2.7.3.2 Chromium 21

2.7.3.3 Sulphur 21

2.7.3.4 Copper 21

2.7.3.5 Aluminium 22

2.7.3.6 Cadmium 22

2.7.3.7 Zinc 23

2.7.3.8 Lead 23

2.7.3.9 Microbial Influences 24

2.7.3.10 Role of Sludge Organic Matter 25

2.7.3.11 Role of sludge iron and manganese 26

CHAPTER 3: EXTENT OF DEDICATED LAND DISPOSAL IN SOUTH AFRICA 27

3.1 Background Information 27

3.1.1 Telephonic survey 27

3.2 Survey of 100 selected sites 29

3.2.1 Macronutrient content of soil samples 31

3.2.2 pH(H_2O) of soil samples 32

3.2.3 Clay content of soil samples 34

3.2.4 Organic carbon content of soil samples 35

3.2.5 Heavy metal content of soil samples 36

3.2.5.1 Cadmium 37

3.2.5.2 Cobalt 38

3.5.2.3 Chromium 39

3.5.2.4 Copper 41

3.5.2.5 Nickel 42

3.5.2.6 Lead 43
3.5.2.7 Zinc 44
3.5.2.8 Other elements 45
3.3 Conclusions from the survey on 100 selected sites 46

CHAPTER 4: CASE STUDIES TO DETERMINE MOBILITY OF SLUDGE CONSTITUENTS AT 40 SELECTED DISPOSAL SITES 47

4.1 Background to Site selection and sampling 47
4.2 Background to Analytical Methods 49
4.3 Correlations between analytical methods 51
4.4 Macronutrients and other elements in soil samples of 40 selected sites 53
4.5 Metals in the sludge and soil samples of 40 selected sites 55
4.6 Mobility of metals in soil samples of 40 selected sites 58
4.6.1 Stockpile sites 59
4.6.2 Instant lawn sites 62
4.6.3 Paddy system sites 64
4.6.4 Irrigation sites 66
4.6.5 Flooded sites 68
4.6.6 Dried application sites 68
4.6.7 Comparison between different types of disposal 71
4.7 Groundwater samples 72
4.8 General observations and concerns 73

CHAPTER 5: MODELLING OF METAL MIGRATION IN SLUDGE AMMENDED SOILS 76

5.1 Description of the Modelling Problem 76
5.1.1 The purpose of the modelling investigation 76
5.1.2 The reference behaviour pattern that is to be modelled 76
5.1.3 How to interpret results 77

5.1.4 How to characterise uncertainty and variability 77

5.1.5 Scenarios to be modelled 77

5.2 Determination of the reference behavior pattern 78

5.2.1 Regression and Correlation Studies 86

5.3 Modelling Method - Operational Philosophy 89

5.4 Model Selection 90

5.5 Model Construction 92

5.6 Model Parameterization 94

5.6.1 Methods 94

5.6.2 Solid phase 94

5.6.2.1 Organic Carbon 95

5.6.2.2 Sulphide 95

5.6.2.3 Hydrous Ferric Oxides (HFO) 96

5.6.3 Liquid phase 96

5.6.3.1 Initial Pore water 96

5.6.3.1.1 Redox status 96

5.6.3.1.2 Sewage plant effluent 97

5.6.3.1.3 Peat water 97

5.6.3.2 Recharge water 97

5.6.3.2.1 Redox status 98

5.6.3.2.2 Rain water 98

5.6.3.2.3 Irrigation water 98

5.7 Model Calibration 98

5.7.1 Reproduction of conceptual models 99

5.7.1.1 Reproduction of known chemistry of the system 99

5.7.1.1.1 Solution Chemistry 99

5.7.1.1.2 Retention mechanism 99

5.7.1.1.3 Influence of biological activity 100

5.7.1.2 Modelling Simulations 101

5.7.2 Reproduction of previous field study results in a similar 107

environment
5.8 Model Verification 108
5.8.1 Batch Extraction with Ammonium Nitrate 113
5.8.2. Batch Extraction with EDTA 113
5.9 Scenario Modelling 114
5.9.1 Scenario 1: Elution of column with dilute salt solution 115
5.9.2 Scenario 2: Influx of acidic water 117
5.9.3 Scenario 3: The effect of maintaining the organic carbon content of the top layer 118
5.9.4 Scenario 4: Effect of adding lime to maintain higher pH 119
5.9.5 Scenario 5: Termination of sludge application 120
5.10 Conclusions from the Modelling Studies 121

CHAPTER 6: CONCLUSIONS AND RECOMMENDATIONS 122
6.1 Extent of DLD practices in South Africa 119
6.2 Case studies 120
6.3 Modelling 123
6.4 Recommendations 124

REFERENCES 128
Appendix A: Detailed Case studies
Appendix B: Soil and water data
Appendix C: General chemical characteristics of soils
Appendix D: Results of Statistical Analyses (Modelling)

LIST OF TABLES

Number	Title	Page
2.5:	Sludge production and disposal methods in various countries	10
2.6a:	Maximum permissible total metal content in soil (mg kg^{-1})	15
2.6b:	Baseline concentration range of metals in South African soils and suggested investigation threshold values for soil (mg kg^{-1}) (from Herselman & Steyn, 2001)	16
2.6c:	Guidelines for NH_4NO_3 extractable metals in soil for groundwater protection (mg kg^{-1}) and water quality guidelines (mg l^{-1})	17
3.2:	Statistics on the type of sludge disposed and industrial effluent received at the 100 selected sites (number of sites)	30
3.2.2:	Average pH(H2O) of the soil samples from 100 selected sites	33
3.2.5.1:	Statistics of total (EPA 3050) Cd in the topsoil of the 100 selected sites (mg kg^{-1})	38
3.5.2.2:	Statistics of total (EPA 3050) Co in the topsoil of the 100 selected sites (mg kg^{-1})	39
3.5.2.3:	Statistics of total (EPA 3050) Cr in the topsoil of the 100 selected sites (mg kg^{-1})	40
3.5.2.4:	Statistics of total (EPA 3050) Cu in the topsoil of the 100 selected sites (mg kg^{-1})	42
3.5.2.5:	Statistics of total (EPA 3050) Ni in the topsoil of the 100 selected sites (mg kg^{-1})	43
3.5.2.6:	Statistics of total (EPA 3050) Pb in the topsoil of the 100 selected sites (mg kg^{-1})	44
3.5.2.7:	Statistics of total (EPA 3050) Zn in the topsoil of the 100 selected sites (mg kg^{-1})	45
3.5.2.8:	Semi-quantitative concentration of other elements in the topsoil of the 100 selected sites (mg kg^{-1})	46

Number	Title	Page
4.1:	Number of sites within some variables for selection of 40 sites for detailed studies	48
4.3:	Correlation matrix of different analytical methods for all metals of all samples, only topsoil samples and only subsoil samples	52
4.4:	Number of sites of each disposal method that fall within guideline ranges for Bray I P, total N and organic C	53
4.5a:	Number of sites with topsoil metal concentrations exceeding the guideline levels set for the different analytical methods	56
4.5b:	Total metal content (*aqua regia*) of sewage sludge at the selected 40 sites (mg kg^{-1})	57
4.6.1:	General information of 19 stockpile sites	59
4.6.2:	General information of 5 instant lawn sites	62
4.6.3:	General information of 4 paddy sites	64
4.6.4:	General information of 7 irrigation sites	66
4.6.5:	General information of 2 flooded sites	68
4.6.6:	General information of 3 dried application sites	68
5.2a:	Summary of speculations of metal solid phases in sludge-amended soils	79
5.2b:	Speculations of operationally defined metal solid phases in sludge-amended soils (McGrath & Cegarra, 1992)	80
5.2c:	Conditional stability constants for metals bound to sewage-amended soils (Sposito *et al.*, 1981)	81
5.2d:	The organic acids in the Mixture Model of Sposito *et al.* (1982), with their acidity constants	82
5.2e:	Conditional constants for metals with sludge-amended soils (Fletcher and Beckett, 1987b)	83
5.2.1a:	Pearson correlation coefficients within soil properties (numbers in bold font are statistically significant)	86

Number	Title	Page
5.2.1b:	Depths to which maxima of metal plumes have traveled, a function of interstitial water pH and organic matter concentration (%)	86
5.2.1c:	Pearson correlation coefficients for soil properties with concentrations of metals extracted by different methods (significant correlations are presented in bold font)	88
5.8a:	NH_4EDTA extraction: Core depths (cm) at which maximum concentrations of metals were found	107
5.8b:	EPA 3050 extraction: Core depths (cm) at which maximum concentrations of metals were found	108
5.8c:	NH_4NO_3 extraction: Core depths (cm) at which maximum concentrations of metals were found	109
5.8d:	Metal loadings and characteristics of the average soil to be simulated	110
5.8.1a:	Percent available metal retained in sludge/soil after extraction with ammonium nitrate solution	110
5.9:	Transport parameters for kinetic column model	112
6.2:	Percentage of samples in each soil layer that exceeded the guideline values set for each analytical method	122

LIST OF FIGURES

Number	Title	Page
2.1:	Waste generation in South Africa (DWAF, 1997)	6
3.1.1:	Disposal options used by 234 different sewage treatment facilities (% of total)	28
3.2a:	Location of the 100 selected sampling sites	29
3.2b:	Disposal options of the 100 sites selected for soil sampling	30
3.2.1a:	Graph and table of % of soil samples within different total N% categories for different disposal methods (both industrial and domestic sludge)	31
3.2.1b:	Graph and table of % of soil samples within different P categories for different disposal methods (both industrial and domestic sludge)	32
3.2.2:	Graph and table of % of soil samples within different pH(H_2O) categories for different disposal methods (both industrial and domestic sludge)	33
3.2.3	Graph and table of % of soil samples within different clay categories for different disposal methods (both industrial and domestic sludge)	34
3.2.4:	Graph and table of % of soil samples within different C% categories for different disposal methods (both industrial and domestic sludge)	35
3.2.5:	Soil samples (%) from 100 sites exceeding the MPL for soils	37
4.3:	Distribution of Bray I P, total N and organic C in the soil profiles of selected sites from each disposal method	
4.6:	Examples of the mobility of metals in the soil profile with the different analytical methods	55
4.6.1:	NH_4EDTA extractable Ni, Zn, Cu and Cd in the soil profiles of 7 selected stockpiled sites (sites 1,2,11,12,15,17 and 19)	58
4.6.2:	NH_4EDTA extractable Ni, Zn, Cu and Cd in the soil profiles of the 5 instant lawn sites	61

Number	Title	Page
4.6.3:	NH_4EDTA extractable Ni, Zn, Cu and Cd in the soil profiles of the 4 sites using paddies as disposal method	63
4.6.4:	NH_4EDTA extractable Ni, Zn, Cu and Cd in the soil profiles of 4 selected irrigation sites (sites 32, 33, 34 and 35)	65
4.6.5:	NH_4EDTA extractable Ni, Zn, Cu and Cd in the soil profiles of the flooded sites	67
4.6.6:	NH_4EDTA extractable Ni, Zn, Cu and Cd in the soil profiles of the dried application sites	69
4.6.7:	Zn concentrations with all analytical methods in the soil profiles of the most contaminated sites from each disposal method	70
4.7:	NO_3 contents of groundwater samples at 9 sites	71
4.8a:	Photographs of two sites situated on soils with low clay content (<5%).	72
4.8b:	Photographs of two sites without fences, one very near a populated area and the other with cattle grazing on the disposal site	74
4.8c:	Photograph of a site situated on a slope near a surface water body without any erosion and run-off control measures	75
4.8d:	Photograph of a site with poor management practices (raw sewage is disposed on disposal site)	75
5.7.1.2a:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution. The soil consists mainly of sand, with hydrous ferric oxide adsorbed to it.	101
5.7.1.2b:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution containing a small concentration of humic acids (pH = 7). The soil consists mainly of sand, with hydrous ferric oxide adsorbed to it.	102

Number	Title	Page
5.7.1.2c:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution containing dilute mineral acid (pH = 3). The soil consists mainly of sand, with hydrous ferric oxide adsorbed to it.	103
5.7.1.2d:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution at approximately neutral pH. The soil consists mainly of sand, with hydrous ferric oxide adsorbed to it.	104
5.7.1.2e:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution at acidic pH. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	104
5.7.1.2f:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution at neutral pH, containing sludge-derived humic acid. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	105
5.8.2:	Elution of a soil column of 20 cells depth by 100 pore volumes of a diammonium EDTA solution. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	111
5.9.1:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute salt solution at neutral pH. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	113
5.9.2a:	Elution of a soil column of 20 cells depth by 100 pore volumes of a dilute acidic solution. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	114

5.9.2b:	Elution of a soil column of 20 cells depth by 100 pore volumes of a near-neutral solution. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	115
5.9.3:	Elution of a soil column of 20 cells depth by 100 pore volumes of a solution in contact with decaying organic matter. The soil consists mainly of sand, with particulate organic matter adsorbed to it.	116
5.9.4:	Elution of a soil column of 20 cells depth by 100 pore volumes of a solution of moderately concentrated lime ([Ca] = 5.0M). The soil consists mainly of sand, with particulate organic matter adsorbed to it.	117