Rainwater harvesting and conservation practices: Challenges and opportunities for sustainable land and water use of "The Green Village" in arid to semi-arid climate zones

Leon van Rensburg & Chris du Preez

27 September 2013







Tillage practices






Tillage practices

Conventional






- Conservation
 - Minimum Tillage
 - Stubble mulch
 - Zero tillage







Tillage practices: 1980-2000

Author	Locality	Ym (kg ha ⁻¹)	Number of
			Seasons
Snyman (1985)	Viljoenskroon	6933-8775	2
	Sannieshof	4063-5160	3
	Delareyville +	3680-4557	2
	Coligny		
Van der Merwe	Sandvet	2595-4948	3
(1993)			
Du Plessis	Bloemfontein	1580-1770	4
(1991)	Bloemfontein	1190-3690	4
	Bloemfontein	320-3640	4
Burger (1982)	Wesselsbron	6273-7082	4
Snyman (1988)	Lichtenburg	690-6756	6







McPhee and Smithen (1985)

% Runoff = 47.9 – 0.44 (% mulch cover)

File name 7-Oct-13

Tillage practices: 1980-2000





□ Scientific contributions


- Bennie, A.T.P., Hoffman, J.E., Coetzee, M.J. & Vrey, H.S., 1994. Storage and utilization of rain water in soils for stabilizing crop production in semi-arid areas [Afr]. Report No 227/1/94. Water Research Commission, Pretoria, South Africa.
- Bennie, A.T.P., Strydom, M.G. & Vrey, H.S., 1998. The application of computer models for agricultural water management on ecotope level [Afr]. Report No. 625/1/98. Water Research Commission, Pretoria, South Africa.
- Bennie, A.T.P. & Hensley, M., 2001. Maximizing precipitation utilization in dryland agriculture in South Africa a review. Journal of Hydrology 241, 124-139.
- ☐ Output: SWAMP model (UFS-WRC)
- ☐ **Impact:** Sustainable decisions on selection of best tillage practices based on soil physical and agronomical principles



Relative soil C loss (du Preez et al., 2011)








Soil C loss from agro-ecosystems (du Preez et al., 2011)








Soil C (%) vs tillage practices (32 yrs) [Loke et al., 2012]







File name

Wheat yield (Mg ha-1)) [Loke et al., 2012]





Period	No-till	Mulch	Plough
1979-2010	2.11	2.19	2.28
1979-1990	2.04	2.07	2.16
1991-2000	1.85	1.90	2.02
2001-2010	2.33	2.29	2.46

Tillage practices: After 2000





- Conventional
- Conservation
 - Minimum Tillage
 - Stubble
 - Zero tillage
 - Rainwater harvesting




HENSLEY, M., BOTHA, J.J. ANDERSON, J.J., VAN STADEN, P.P. & DU TOIT, A., 2000. Optimizing rainfall use efficiency for developing farmers with limited access to irrigation water. Report No. 878/1/00. Water Research Commission, Pretoria, South Africa.

In-field rainwater harvesting: Bio-physical













In-field rainwater harvesting: Mulching






Botha JJ, Van Rensburg LD, Anderson JJ, Hensley M, Macheli MS, Van Staden PP, Kundhlande G, Groenewald DG, Baiphethi MN. 2003. *Water conservation techniques on small plots in semi-arid areas to enhance rainfall use efficiency, food security, and sustainable crop production.* Water Research Commission, Report no 1176/1/03, Pretoria, South Africa.













IRWH: Technology exchange







In-field rainwater harvesting: Socio-economic



Kundhlande G, Groenewald DG, Baiphethi MN, Viljoen MF, Botha JJ, van Rensburg LD, Anderson JJ. 2004. Socio-economic study on water conservation techniques in semi-arid areas. Water Research Commission, Report no 1267/1/04, Pretoria, South Africa.













IRWH-upscaling (van Rensburg et al., 2012)
















Impact





- Number of WRC
- Universities
- Capacity building
- Society
- International





Conclusion: Forgotten green village













File name